

Touching the $\mathbb{Z}_2$ in Three-Dimensional Rotations

Vesna Stojanoska,^{*} Orlin Stoytchev[†]

Abstract

A simple and self-contained proof is presented of the well-known fact that the fundamental group of $SO(3)$ is $\mathbb{Z}_2$, using a relationship between closed paths in $SO(3)$ and braids.

1 Introduction

The three-dimensional rotations form a Lie group, usually denoted by $SO(3)$. A fascinating feature of our world is that the fundamental group of $SO(3)$ is $\mathbb{Z}_2$, or in short $\pi_1(SO(3)) \cong \mathbb{Z}_2$. This means that all closed paths in $SO(3)$ starting and ending at the identity fall into two homotopy classes — those that are homotopic to the constant path and those that are not. Composing two paths from the second class yields a path from the first class. This topological property of the space parametrizing rotations can also be stated as follows — a complete rotation of an object (corresponding to a closed path in $SO(3)$) may or may not be continuously deformable to the trivial motion (i.e., no motion at all); the composition of two motions that are not deformable to the trivial one gives a motion which is.

A manifold with a fundamental group $\mathbb{Z}_2$ is a challenge to our imagination — it is easy to visualize spaces with fundamental group $\mathbb{Z}$, (the punctured plane), or $\mathbb{Z} \times \mathbb{Z} \cdots \times \mathbb{Z}$, (plane with several punctures), or even $\mathbb{Z} \oplus \mathbb{Z}$ (torus), but there is no two-dimensional manifold embedded in $\mathbb{R}^3$ whose fundamental group is $\mathbb{Z}_2$.

The peculiar structure of $SO(3)$ plays a crucial role in our physical world. We know that there are two quite different types of elementary particles, bosons and fermions. The quantum state of a boson is described by a (possibly multi-component) wave function, which remains unchanged when a full (360°) rotation of the coordinate system is performed, while the wave function of a fermion gets multiplied by -1 under a complete rotation. Somewhat loosely speaking, this comes from the fact that only the modulus of the wave function has a direct physical meaning and therefore the wave function need not transform under a true representation of $SO(3)$ but just under a projective representation, which is a true representation of its universal covering group $SU(2)$ [6, 2].

The standard way of showing that $\pi_1(SO(3)) \cong \mathbb{Z}_2$ is to prove that $SO(3)$ is homeomorphic to S^3/i , where i denotes the identification of diametrically opposite points of S^3 . Once the latter is known, it is quite easy to see that paths starting from one pole of S^3 and ending at the other pole will be closed paths in S^3/i , which are not contractible, but the composition of two such paths gives a contractible path. In order to find the topological structure of $SO(3)$ one normally uses Lie group and Lie algebra theory. Namely, it is shown that the group $SU(2)$ of unitary 2×2 matrices with determinant 1 is homeomorphic to S^3 and that there is a 2–1 homomorphism $SU(2) \rightarrow SO(3)$,

^{*}American University in Bulgaria, 2700 Blagoevgrad, Bulgaria, vstotka@gmail.com, vns021@aubg.bg

[†]American University in Bulgaria, 2700 Blagoevgrad, Bulgaria and Institute for Nuclear Research, 1784 Sofia, Bulgaria, ostroytchev@aubg.bg

which is a local isomorphism, and which sends diametrically opposite points in $SU(2)$ to the same point in $SO(3)$.

There is a more direct way to exhibit the topological structure of the parameter space of three-dimensional rotations. It only uses the fact that rotations are represented by 3×3 matrices and any such matrix must have one (real) eigenvalue and a corresponding eigenvector. Then one shows easily that this element of $SO(3)$ must be a rotation around this eigenvector. In other words any element of $SO(3)$ is a rotation around some axis and we need to specify the angle of rotation and the orientation of that axis in $\mathbb{R}^3$.

In the present paper we present an alternative way of proving that $\pi_1(SO(3)) \cong \mathbb{Z}_2$. It does not use Lie group theory or even matrices. It is purely algebraic-topological in nature and very visual. It displays a simple connection between full rotations (closed paths in $SO(3)$) and braids. We believe that this may be an interesting way of showing a nontrivial topological result to students in introductory geometry and topology courses as well as a suitable way of introducing braids and braid groups.

The goal of this paper is mostly pedagogical — presenting in a self-contained and accessible way a set of results that are basically known to algebraic topologists and people studying braid groups. The fact that the first homotopy group of $SO(3)$ can be related to spherical braids is a special case (in disguise) of the following general statement [4]: “The configuration space of three points on an r -sphere is homotopically equivalent to the Stiefel manifold of orthogonal two-frames in $r + 1$ -dimensional Euclidean space”. Fadell [4] considers a particular element of $\pi_1(SO(3))$ and uses the fact that it has order 2 to prove a similar statement for a corresponding braid. Our direction is opposite - we analyze braids to deduce topological properties of $SO(3)$.

In the next section we describe a simple experiment that actually demonstrates the $\mathbb{Z}_2$ in three-dimensional rotations. Then in section 3 we give a formal treatment of that experiment. We construct a map from $\pi_1(SO(3))$ into a certain factorgroup of a subgroup of the braid group with three strands. We prove that this map is an isomorphism and that the image is $\mathbb{Z}_2$.

2 The experiment

Take a ball (tennis ball will do) and attach three strands to three different points on its surface. Attach the other ends of the strands to three different points on the surface of your desk (Figure 1). Perform an arbitrary number of full rotations of the ball around arbitrary axes. You will get an



Figure 1: Rotating a ball with strands attached

entangled “braid”. Now keep the orientation of the ball fixed. If the total number of full rotations is even, you can always untangle the “braid” by flipping strands around the ball. If the number of rotations is odd you will never be able to untangle it, but you can always reduce it to one simple

configuration, e.g., the one obtained by rotating the ball around the first point and twisting the second and third strands around each other.

As is to be expected, rotations that can be continuously deformed to the trivial rotation (i.e., no rotation) lead to trivial braiding. At this point we can only conjecture from our experiment that the fundamental group of $SO(3)$ contains as a factor $\mathbb{Z}_2$.

3 Relating three-dimensional rotations to braids

With each closed path in $SO(3)$ we associate three closed paths in $\mathbb{R}^3$ starting at the sphere with radius 1 and ending at the sphere with radius $1/2$. We may think of continuously rotating a sphere from time $t = 0$ to time $t = 1$ so that the sphere ends up with the same orientation as the initial one. Simultaneously we shrink the radius of the sphere from 1 to $1/2$ (see Figure 2). Any three points on




Figure 2: A “spherical braid” and a normal braid

the sphere will trace three continuous paths in $\mathbb{R}^3$, which do not intersect each other. Furthermore, for fixed t the three points on these paths lie on the sphere with radius $1 - t/2$. To formalize things, let $\omega(t)$, $t \in [0, 1]$ be any continuous path in $SO(3)$ with $\omega(0) = \omega(1) = I$. $\omega(t)$ acts on vectors (points) in $\mathbb{R}^3$. Take three initial points in $\mathbb{R}^3$, e.g., $\mathbf{x}_0^1 = (1, 0, 0)$, $\mathbf{x}_0^2 = (-1/2, \sqrt{3}/2, 0)$, $\mathbf{x}_0^3 = (-1/2, -\sqrt{3}/2, 0)$. Define three continuous paths by

$$\mathbf{x}^i(t) := (1 - t/2)\omega(t)(\mathbf{x}_0^i), \quad t \in [0, 1], \quad i = 1, 2, 3 .$$

In this way we get an object that will be called a *spherical braid* — several distinct points on a sphere and the same number of points, in the same positions, on a smaller sphere, connected by strands in such a way that the radial coordinate of each strand is monotonic in t .

Note One can multiply two spherical braids by connecting the ends of the first to the beginnings of the second (and rescaling the parameter). When one considers classes of isotopic spherical braids one obtains the so called *braid group of the sphere* [5], which algebraically is B_3/R (see below). This is known as the mapping-class group of the sphere (with 0 punctures and 0 boundaries) and has been studied by topologists.

We can map our spherical braid to a conventional one using stereographic projection (Figure 2). First we choose a ray starting at the origin and not intersecting any strand. Then we map stereographically any point on a sphere with radius $1/2 \leq \rho \leq 1$, except the point where the ray intersects that sphere and with respect to which we project, to a point in a corresponding horizontal plane. Finally we define the z -coordinate of the image to be $z = -\rho$.

Recall the usual notion of *braids*, introduced by Artin [1]. (See also [3] for a contemporary review of the theory of braids and its relations to other subjects.) One takes two planes in $\mathbb{R}^3$, let's say parallel to the XY plane, fixes n distinct points on each plane and connects each point on the lower plane with a point on the upper plane by a continuous path (strand). The strands do not intersect each other. In addition the z -coordinate of each strand is a monotonic function of



Figure 3: The generators σ_1 and σ_2 of B_3




Figure 4: The braid relation for B_3

the parameter of the strand and thus z can be used as a common parameter for all strands. Two different braids are considered equivalent or *isotopic* if there exists a homotopy of the strands (keeping the endpoints fixed), so that for each value of the homotopy parameter s you get a braid, for $s = 0$ you get the initial braid and for $s = 1$ the final one. When the points on the lower and the upper plane have the same positions (their x and y coordinates are the same), we can multiply braids by stacking one on top of the other. Considering classes of isotopic braids with the multiplication just defined, the *braid group* is obtained. Artin showed that the braid group B_n on n strands has a presentation with $n - 1$ generators and a simple set of relations - Artin's braid relations. We give them for the case $n = 3$ since this is the one we are mostly interested in. In this case the braid group B_3 is generated by the generators σ_1 , corresponding to twisting of the first and the second strands, and σ_2 , corresponding to twisting of the second and the third strands (the one to the left always passing behind the one to the right) (Figure 3). These generators are subject to a single braid relation (Figure 4):

$$\sigma_2 \sigma_1 \sigma_2 = \sigma_1 \sigma_2 \sigma_1 \quad (1)$$

We say that B_3 has a *presentation* with generators σ_1 and σ_2 and defining relation given by Equation 1, or in short:

$$B_3 = \langle \sigma_1, \sigma_2; \sigma_1 \sigma_2 \sigma_1 \sigma_2^{-1} \sigma_1^{-1} \sigma_2^{-1} \rangle \quad (2)$$

In our case, since a full rotation of the sphere returns the three points to their original positions, we always get *pure braids*, i.e., braids for which any strand connects a point on the lower plane with its translate on the upper plane. Pure braids form a subgroup of B_3 which is denoted by P_3 . Note that intuitively there is a homomorphism π from B_3 to the symmetric group S_3 since any braid from B_3 permutes the three points. Formally one defines π on the generators by

$$\pi(\sigma_1)(1, 2, 3) = (2, 1, 3), \quad \pi(\sigma_2)(1, 2, 3) = (1, 3, 2) \quad (3)$$

and then extends it to the whole group B_3 (it is important that π maps Equation 1 to the trivial identity). Pure braids are precisely those that do not permute the points and therefore we can give

the following algebraic characterization of P_3 :

$$P_3 := \text{Ker } \pi$$

Alternatively, S_3 is the quotient of B_3 by the additional equivalence relations $\sigma_i^2 \sim I$, $i = 1, 2$ and if N is the minimal normal subgroup containing σ_i^2 , then $\pi : B_3 \rightarrow B_3/N$ is the natural projection. It is then easy to see that the kernel of π has to be a product of words of the following type:

$$\sigma_{i_1}^{\pm 1} \sigma_{i_2}^{\pm 1} \dots \sigma_{i_k}^{\pm 1} \sigma_{i_{k+1}}^{\pm 2} \sigma_{i_k}^{\pm 1} \dots \sigma_{i_2}^{\pm 1} \sigma_{i_1}^{\pm 1}$$

The whole subgroup P_3 can in fact be generated by the following three *twists* (Figure 5)




Figure 5: The generators a_{12} , a_{13} and a_{23} of P_3

$$a_{12} := \sigma_1^2, \quad a_{13} := \sigma_2 \sigma_1^2 \sigma_2^{-1} = \sigma_1^{-1} \sigma_2^2 \sigma_1, \quad a_{23} := \sigma_2^2 \quad (4)$$

In our construction so far we mapped any closed path in $SO(3)$ to a spherical braid and then, using stereographic projection, to a conventional pure braid. The last map, however, depends on a choice of a ray in $\mathbb{R}^3$ and, what is worse, spherical braids that are isotopic (in the obvious sense) may map to nonisotopic braids. To mend this, we will identify certain classes of braids in P_3 . Namely, we introduce the following equivalence relations (see Figure 6):




Figure 6: The flips r_1 , r_2 and r_3

$$r_1 := \sigma_1 \sigma_2^2 \sigma_1 \sim I, \quad r_2 := \sigma_1^2 \sigma_2^2 \sim I, \quad r_3 := \sigma_2 \sigma_1^2 \sigma_2 \sim I. \quad (5)$$

On our model with the tennis ball from Section 2 r_i , $i = 1, 2, 3$ correspond to *flips* of the i -th strand above and around the ball. Such motions lead to isotopic spherical braids, as will be shown later.

Note Strictly speaking, we have to make sure that our final result will not change when any strand crosses at any part of the braid the ray which we use for the stereographic projection. This means that we have to factorize by the normal closure in B_3 (not in P_3 !) of the generators r_i , $i = 1, 2, 3$,

i.e., the smallest normal subgroup in B_3 containing these three generators. This would then allow us to set to I any r_i in any part of a word. One sees immediately that only one of the generators is needed then, since the other two will be contained in the normal closure of the first. We noticed, however, that we managed to untie any trivial braid just by a sequence of the three flips r_i defined in Equation 5 and their inverses, performed at the end of the braid. At the same time a nontrivial braid, corresponding to an odd number of rotations, cannot be untied even if we allow flips in any part of the braid. This can only be true if the flips r_i generate a normal subgroup in B_3 (which of course then coincides with the normal closure of any of the r_i and is also normal in P_3).

Lemma 1

The subgroup $R \subset P_3$, generated by r_1, r_2, r_3 is normal in B_3 .

Proof: We need to show that we can represent all conjugates of r_i with respect to the generators of B_3 and their inverses as products of the r_i and their inverses. Straightforward calculations, using multiple times Artin's braid relation (Equation 1) give the following identities:

$$\begin{aligned} \sigma_1 r_1 \sigma_1^{-1} &= r_2, & \sigma_2 r_1 \sigma_2^{-1} &= \sigma_2^{-1} r_1 \sigma_2 = r_1, \\ \sigma_1 r_2 \sigma_1^{-1} &= r_2 r_1 r_2^{-1}, & \sigma_2 r_2 \sigma_2^{-1} &= r_3, \\ \sigma_1 r_3 \sigma_1^{-1} &= \sigma_1^{-1} r_3 \sigma_1 = r_3, & \sigma_2 r_3 \sigma_2^{-1} &= r_1^{-1} r_2 r_1 = r_3 r_2 r_3^{-1}, \\ \sigma_1^{-1} r_1 \sigma_1 &= r_1^{-1} r_2 r_1, & \sigma_1^{-1} r_2 \sigma_1 &= r_1, \\ \sigma_2^{-1} r_2 \sigma_2 &= r_1 r_3 r_1^{-1} = r_2^{-1} r_3 r_2, & \sigma_2^{-1} r_3 \sigma_2 &= r_2. \end{aligned}$$

We demonstrate as an example the proof of the first identity in the second line. We have

$$\begin{aligned} \sigma_1 \sigma_2 \sigma_1 &= \sigma_2 \sigma_1 \sigma_2 \\ \sigma_2 \sigma_1 \sigma_2 \sigma_1 \sigma_2 \sigma_1 &= \sigma_2^2 \sigma_1 \sigma_2^2 \sigma_1 \\ \sigma_1 \sigma_2 \sigma_1 \sigma_2 \sigma_1 \sigma_2 &= \sigma_2^2 \sigma_1 \sigma_2^2 \sigma_1 \\ \sigma_1 \sigma_2^2 \sigma_1 \sigma_2^2 &= \sigma_2^2 \sigma_1 \sigma_2^2 \sigma_1 \\ \sigma_1 \sigma_2^2 \sigma_1 &= \sigma_2^2 \sigma_1 \sigma_2^2 \sigma_1 \sigma_2^{-2} \\ \sigma_1^3 \sigma_2^2 \sigma_1^{-1} &= \sigma_1^2 \sigma_2^2 \sigma_1 \sigma_2^2 \sigma_1 \sigma_2^{-2} \sigma_1^{-2} \end{aligned}$$

and therefore

$$\sigma_1 r_2 \sigma_1^{-1} = \sigma_1 \cdot \sigma_1^2 \sigma_2^2 \cdot \sigma_1^{-1} = \sigma_1^2 \sigma_2^2 \cdot \sigma_1 \sigma_2^2 \sigma_1 \cdot \sigma_2^{-2} \sigma_1^{-2} = r_2 r_1 r_2^{-1}$$

By suitable full rotations we obtain all generators of P_3 . For example, a_{12} is obtained by rotating around the vector $\mathbf{x}_0^3 = (-1/2, -\sqrt{3}/2, 0)$ and it twists the first and the second strand. Furthermore, homotopies between closed paths in $SO(3)$ correspond to isotopies of the spherical braids and thus homotopic closed paths in $SO(3)$ will be mapped to the same element in the factorgroup P_3/R .

Proposition 1

The factorgroup P_3/R is isomorphic to $\mathbb{Z}_2$.

Proof: To make notations simpler we use the same letter to denote both a representative of a class in P_3/R and the class itself, hoping that the meaning is clear from the context. In P_3/R we have

$$\sigma_1 \sigma_2^2 = \sigma_1^{-1} = \sigma_2^2 \sigma_1,$$

and

$$\sigma_2 \sigma_1^2 = \sigma_2^{-1} = \sigma_1^2 \sigma_2.$$

Now we obtain the following sequence of identities:

$$\begin{aligned}\sigma_2\sigma_1^2 &= \sigma_1^2\sigma_2, & \sigma_1\sigma_2\sigma_1^2 &= \sigma_1^3\sigma_2, \\ \sigma_2\sigma_1\sigma_2\sigma_1 &= \sigma_1^3\sigma_2, & \sigma_1\sigma_2\sigma_1\sigma_2\sigma_1 &= \sigma_1^4\sigma_2, \\ \sigma_1\sigma_2^2\sigma_1\sigma_2 &= \sigma_1^4\sigma_2, & I &= \sigma_1^4.\end{aligned}$$

We have used twice the braid relation (Equation 1) and the first equivalence relation in Equation 5. In completely the same way we prove

$$\sigma_2^4 = I.$$

Combining the last two results with the equivalence relations (Equation 5) we finally get

$$\sigma_1^2 = \sigma_1^{-2} = \sigma_2^2 = \sigma_2^{-2} \quad (6)$$

It is now clear that in P_3/R the three generators, defined in Equation 4 reduce to one element of order 2. Therefore they generate $\mathbb{Z}_2$. This completes the proof.

So far we have constructed a map $\pi_1(SO(3)) \rightarrow P_3/R$, which is onto by construction, and we have shown that the image is isomorphic to $\mathbb{Z}_2$. We now show that this map is actually an isomorphism.

Proposition 2

The map $\pi_1(SO(3)) \rightarrow P_3/R$ is a monomorphism.

Proof: We have to show that if a closed continuous path in $SO(3)$ is mapped to a braid in R , then this path is homotopic to the constant path. The proof basically reduces to the following observation — any spherical braid which is pure (the strands connect each point on the outer sphere with the same point on the inner sphere) determines a closed path in $SO(3)$. Two isotopic spherical pure braids determine homotopic closed paths in $SO(3)$. Indeed, recall that for a spherical braid we can parametrize the points on each strand with a single parameter t and that for a fixed t all three points lie on a sphere with radius $1 - t/2$. These three ordered points $\mathbf{x}^i(t), i = 1, 2, 3$ give for every fixed t a nondegenerate triangle, oriented somehow in $\mathbb{R}^3$. Let $\mathbf{l}(t)$ be the vector, connecting the center of mass of the triangle with the vertex $\mathbf{x}^1(t)$, i.e., $\mathbf{l}(t) = \mathbf{x}^1 - (\mathbf{x}^1(t) + \mathbf{x}^2(t) + \mathbf{x}^3(t))/3$ and define $\mathbf{e}^1(t) := \mathbf{l}(t)/\|\mathbf{l}(t)\|$. Let $\mathbf{e}^3(t)$ be the unit vector, perpendicular to the plane of the triangle, in a positive direction relative to the orientation (1,2,3) of the boundary. Finally, let $\mathbf{e}^2(t)$ be the unit vector, perpendicular to both $\mathbf{e}^1(t)$ and $\mathbf{e}^3(t)$, so that the three form a right-handed frame. Then there is a unique element $\omega(t) \in SO(3)$ sending the vectors $\mathbf{e}_0^1 = (1, 0, 0)$, $\mathbf{e}_0^2 = (0, 1, 0)$, $\mathbf{e}_0^3 = (0, 0, 1)$ to the triple $\mathbf{e}^i(t)$. According to our definitions, $\omega(0) = \omega(1) = I$ and we get a continuous function $\omega : [0, 1] \rightarrow SO(3)$, where continuity should be understood relative to some natural topology on $SO(3)$, e.g., the strong operator topology.

Now, if we have two isotopic spherical braids, by definition there are continuous functions $\mathbf{x}^i(t, s), i = 1, 2, 3$ such that $\mathbf{x}^i(t, s)$ is a braid for any fixed $s \in [0, 1]$, $\mathbf{x}^i(0, s) = \mathbf{x}_0^i$, $\mathbf{x}^i(1, s) = \mathbf{x}_0^i/2$, $\mathbf{x}^i(t, 0)$ give the initial braid and $\mathbf{x}^i(t, 1)$ give the final braid. By assigning an element $\omega(t, s)$ to any triple $\mathbf{x}^i(t, s)$ as described we get a homotopy between two closed paths in $SO(3)$.

Let $\omega'(t)$ be a closed path in $SO(3)$ which is mapped to a braid b in the class $r_1 \in R$. We can construct a spherical braid, whose image is isotopic to that braid. Let $\mathbf{z}$ be the point on the unit sphere with respect to which we perform the stereographic projection. This can always be chosen to be the north pole or a point very close to the north pole (in case a strand is actually crossing the axis passing through the north pole). Note that the points $\mathbf{x}_0^i, i = 1, 2, 3$ are on the equator. Construct a simple closed path on the unit sphere starting and ending at $\mathbf{x}_0^1$ and going around $\mathbf{z}$ in a negative direction (without crossing the equator except at the endpoints). Thus we have two continuous functions $\varphi(t), \theta(t), t \in [0, 1]$ — the spherical (angular) coordinates describing this path. Let $\mathbf{x}^1(t)$ be the point in $\mathbb{R}^3$ whose spherical coordinates are $\rho(t) := 1 - t/2$, $\varphi(t)$, $\theta(t)$ and

let $\mathbf{x}^i(t) := (1 - t/2)\mathbf{x}_0^i, i = 2, 3$. These three paths give the required spherical braid. It is isotopic to the trivial braid, coming from the constant path in $SO(3)$ and at the same time it is isotopic to the preimage of b under the stereographic projection. In this way we see that $\omega'(t)$ must be homotopic to the constant path. Obviously similar argument holds with r_1 replaced by r_2 and r_3 or the inverses. Since any element in R is a product of these generators and since products of isotopic braids give isotopic braids, this completes the proof.

4 Further results and generalizations

There is an obvious generalization of some of the results of the previous sections to the case $n > 3$. The minimal number of strands that is needed to capture the nontrivial fundamental group of $SO(3)$ is $n = 3$. When $n > 3$ any full rotation will give rise to a pure spherical braid but the whole group of pure braids will not be generated in this way. It is relatively easy to see that in this way, after projecting stereographically, one will obtain a subgroup of P_n , generated by a single *full twist* d of all strands around an external point and a set of n flips r_i :

$$\begin{aligned} d &:= (\sigma_{n-1} \cdots \sigma_2 \sigma_1)^n, \\ r_1 &:= \sigma_1 \sigma_2 \cdots \sigma_{n-2} \sigma_{n-1}^2 \sigma_{n-2} \cdots \sigma_1, \\ r_2 &:= \sigma_1^2 \sigma_2 \cdots \sigma_{n-2} \sigma_{n-1}^2 \sigma_{n-2} \cdots \sigma_2, \\ r_i &:= \sigma_{i-1} \cdots \sigma_2 \sigma_1^2 \sigma_2 \cdots \sigma_{n-2} \sigma_{n-1}^2 \sigma_{n-2} \cdots \sigma_i, \quad i = 2, 3, \dots, n-1, \\ r_n &:= \sigma_{n-1} \sigma_{n-2} \cdots \sigma_2 \sigma_1^2 \sigma_2 \cdots \sigma_{n-2} \sigma_{n-1}. \end{aligned}$$

Figure 7 shows a full twist for the case with 3 strands while Figure 8 shows a generic flip.



Figure 7: The full twist d in the case $n = 3$



Figure 8: The flip r_i

Straightforward calculations give the following generalization of Lemma 1:

Lemma 1'

The subgroup $R \subset P_n$, generated by r_i , $i = 1, \dots, n$, is normal in B_n .

Proof: As in the proof of Lemma 1 we exhibit explicit formulas for the conjugates of all flips r_i :

$$\begin{aligned}\sigma_j r_i \sigma_j^{-1} &= \sigma_j^{-1} r_i \sigma_j = r_i, \quad i - j > 1 \text{ or } j - i > 0, \\ \sigma_{i-1} r_i \sigma_{i-1}^{-1} &= r_i r_{i-1} r_i^{-1}, \\ \sigma_{i-1}^{-1} r_i \sigma_{i-1} &= r_{i-1}, \\ \sigma_i r_i \sigma_i^{-1} &= r_{i+1}, \quad i < n - 1 \\ \sigma_i^{-1} r_i \sigma_i &= r_i^{-1} r_{i+1} r_i, \quad i < n - 1\end{aligned}$$

Let us denote by S the subgroup, generated by d and r_i . Using purely topological information, namely that $\pi_1(SO(3)) \cong \mathbb{Z}_2$, we can deduce the following generalization of Proposition 1:

Proposition 1'

The factorgroup S/R is isomorphic to $\mathbb{Z}_2$.

An equivalent statement is that $d^2 \in R$.

It is tempting to try generalizing the main result of this paper to higher-dimensional rotations. A straightforward generalization fails since one would produce braids in higher than three-dimensional space and these can always be untangled. Note that in three dimensions we are able to attribute a path in $SO(3)$ to any spherical braid with 3 strands but this is not the case for $n > 3$ (4 points on S^3 may not determine an orientation of the orthonormal frame in $\mathbb{R}^4$.) It remains to be seen if a refinement of the methods used can render meaningful results.

References

- [1] E. Artin, *Theory of Braids*, Math. Ann. **48**, No. 1 (1947), 101–126.
- [2] V. Bargmann, *On Unitary Ray Representations of Continuous Groups*, Math. Ann. **59**, No. 1 (1954), 1–46.
- [3] J. Birman and T. Brendle, *Braids: a Survey*, E-print math.GT/0409205, <http://arxiv.org/abs/math/0409205>.
- [4] E. Fadell, *Homotopy Groups of Configuration Spaces and the String Problem of Dirac*, Duke Math. J. **29** (1962), 231–242.
- [5] E. Fadell and J. Van Buskirk, *The Braid Groups of E^2 and S^2* , Duke Math. J. **29** (1962), 243–257.
- [6] E. Wigner, *On Unitary Representations of the Inhomogeneous Lorentz Group*, Math. Ann. **40**, No. 1 (1939), 149–204.